



Do uncontrollable movements
make you feel like

disconnectING?

It could be tardive dyskinesia (TD).

Ask your healthcare provider about
reduc**ING** your TD movements with
INGREZZA® (valbenazine) capsules.



Learn more at
[INGREZZA.com/TD](https://www.ingrezza.com/TD)

Actor portrayals



Important Information

Approved Use

INGREZZA® (valbenazine) capsules or INGREZZA® SPRINKLE (valbenazine) capsules are prescription medicines used to treat adults with movements in the face, tongue, or other body parts that cannot be controlled (tardive dyskinesia).

It is not known if INGREZZA or INGREZZA SPRINKLE is safe and effective in children.

SELECT IMPORTANT SAFETY INFORMATION

INGREZZA or INGREZZA SPRINKLE can cause serious side effects in people with Huntington's disease, including: depression, suicidal thoughts, or suicidal actions. Tell your healthcare provider before you start taking INGREZZA or INGREZZA SPRINKLE if you have Huntington's disease and are depressed (have untreated depression or depression that is not well controlled by medicine) or have suicidal thoughts. Pay close attention to any changes, especially sudden changes, in mood, behaviors, thoughts, or feelings. This is especially important when INGREZZA or INGREZZA SPRINKLE is started and when the dose is changed. Call your healthcare provider right away if you become depressed, have unusual changes in mood or behavior, or have thoughts of hurting yourself.

Please see additional Important Safety Information on pages 18–19 and accompanying full **[Prescribing Information](#)**, including Boxed Warning, and **[Medication Guide](#)**.

Tardive dyskinesia: Persistent, uncontrollable body movements

If you have persistent, uncontrollable movements and have taken certain mental health meds, it's time to ask your healthcare provider about a treatable condition called tardive dyskinesia (TD).

TD movements can occur anywhere in the body:

UPPER BODY

ARMS, TORSO,
HANDS/FINGERS

LOWER BODY

LEGS,
FEET/TOES



FACE

MOUTH, LIPS,
TONGUE, JAW,
EYES

Take an important step:
Understand the cause of your
uncontrollable movements and
learn about treatment.

Visit [INGREZZA.com/Movements](https://www.ingrezza.com/Movements)

What medicines can cause TD?

You may have been prescribed certain mental health medicines (antipsychotics) to treat a condition such as:

Depression

Schizophrenia/schizoaffective disorder

Bipolar disorder

Anxiety disorder

Certain prescription medicines used to treat gastrointestinal disorders (metoclopramide and prochlorperazine) may also cause TD.

If you are currently taking antipsychotic medicine, do not stop, adjust, or switch your medicine without talking to your healthcare provider.

View a list of meds known to cause TD at [INGREZZA.com/Meds-to-Monitor](https://www.ingrezza.com/Meds-to-Monitor)

Select Important Safety Information

Do not take INGREZZA or INGREZZA SPRINKLE if you:

- are allergic to valbenazine, or any of the ingredients in INGREZZA or INGREZZA SPRINKLE.

Please see additional Important Safety Information on pages 18–19 and accompanying full [Prescribing Information](#), including Boxed Warning, and [Medication Guide](#).

ONCE-DAILY
INGREZZA[®]
(valbenazine) capsules

Living with tardive dyskinesia

TD is unlikely to get better on its own, but you can take control by talking to your healthcare provider right away.

Even mild TD movements can negatively affect your work, social outings, and daily activities.* Uncontrollable movements can create an unintended impression on the people you meet.

Those movements can also affect your emotional well-being.*



“Because of my TD, I was embarrassed to be out in public. I would chew gum a lot to mask the movement of my jaws. I wouldn’t attend work outings, family functions. I missed a lot of milestones because I would seclude myself.”

– Steve, living with bipolar disorder and tardive dyskinesia

Steve was compensated by Neurocrine Biosciences, Inc. to share his story



Watch real patient stories
at [INGREZZA.com/Stories](https://www.ingrezza.com/Stories)

Is TD impactING you? Ask your healthcare provider about reducing uncontrollable movements with INGREZZA® (valbenazine) capsules—it’s a treatment proven to reduce TD that’s always one capsule, once daily, and #1 prescribed.

*In a survey, 127 people diagnosed with TD were asked, “TD may impact you in many different ways. To what extent has TD impacted you in each of the following areas?” Answers were based on a scale of 1 (not impacted at all) to 7 (extremely impacted).

Select Important Safety Information

INGREZZA or INGREZZA SPRINKLE can cause serious side effects, including:

- **Allergic reactions.** Allergic reactions, including an allergic reaction that causes sudden swelling called angioedema, can happen after taking the first dose or after many doses of INGREZZA or INGREZZA SPRINKLE. Signs and symptoms of allergic reactions and angioedema include: trouble breathing or shortness of breath, swelling of your face, lips, eyelids, tongue, or throat, or other areas of your skin, trouble with swallowing, or rash, including raised, itchy red areas on your skin (hives). Swelling in the throat can be life-threatening and can lead to death. Stop taking INGREZZA or INGREZZA SPRINKLE and go to the nearest emergency room right away if you develop these signs and symptoms of allergic reactions and angioedema.

Please see additional Important Safety Information on pages 18–19 and accompanying full [Prescribing Information](#), including Boxed Warning, and [Medication Guide](#).

ONCE-DAILY
INGREZZA®
(valbenazine) capsules

Car^{ING} for someone with tardive dyskinesia

It can be tough to tell a loved one that you've noticed they have uncontrollable movements. But the sooner they're aware, the sooner they can take control by talking to a healthcare provider about diagnosis and treatment.



“I’m glad I said something to Amy about her uncontrollable movements, because even though she was doing it, she had no clue she was doing it. If we didn’t discuss it, we never would’ve made it to where we are now, and learned about INGREZZA.”

- Kim (right), care partner of Amy, who lives with depression and TD

Amy & Kim were compensated by Neurocrine Biosciences, Inc. to share their stories

“If you have uncontrollable movements, and you’ve had a history of taking medicines for mental health issues, you might have TD. Go to your doctor and advocate for yourself. There are treatments.”

- Forrest (right), care partner of Moira, who lives with depression and TD

Moira & Forrest were compensated by Neurocrine Biosciences, Inc. to share their stories



If your loved one is having uncontrollable movements, they may need your support. It’s OK to speak up. Encourage them to talk to a healthcare provider about TD.

Select Important Safety Information

INGREZZA or INGREZZA SPRINKLE can cause serious side effects, including:

- **Sleepiness and tiredness that could cause slow reaction times (somnolence and sedation).** Do not drive a car or operate dangerous machinery until you know how INGREZZA or INGREZZA SPRINKLE affects you. Drinking alcohol and taking other medicines may also cause sleepiness during treatment with INGREZZA or INGREZZA SPRINKLE.

Please see additional Important Safety Information on pages 18–19 and accompanying full [Prescribing Information](#), including Boxed Warning, and [Medication Guide](#).

ONCE-DAILY
INGREZZA[®]
(valbenazine) capsules

Select Important Safety Information

INGREZZA or INGREZZA SPRINKLE can cause serious side effects, including:

- **Heart rhythm problems (QT prolongation).** INGREZZA or INGREZZA SPRINKLE may cause a heart rhythm problem known as QT prolongation. You have a higher chance of getting QT prolongation if you also take certain other medicines during treatment with INGREZZA or INGREZZA SPRINKLE. Tell your healthcare provider right away if you develop any signs or symptoms of QT prolongation, including: fast, slow, or irregular heartbeat (heart palpitations), shortness of breath, dizziness or lightheadedness, or fainting or feeling like you are going to faint.
- **Neuroleptic Malignant Syndrome (NMS).** NMS is a serious condition that can lead to death. Call a healthcare provider right away or go to the nearest emergency room if you develop these symptoms and they do not have another obvious cause: high fever, stiff muscles, problems thinking, irregular pulse or blood pressure, increased sweating, or very fast or uneven heartbeat.
- **Parkinson-like symptoms.** Symptoms include: body stiffness, drooling, trouble moving or walking, trouble keeping your balance, shaking (tremors), or falls.

Please see additional Important Safety Information on pages 18–19 and accompanying full [Prescribing Information](#), including Boxed Warning, and [Medication Guide](#).

ONCE-DAILY
 **INGREZZA**[®]
(valbenazine) capsules

Treat **ING** TD with INGREZZA



INGREZZA is proven to reduce TD,
always one capsule, once daily,
and #1 prescribed.

INGREZZA® (valbenazine) capsules is used to treat adults with TD in the face, tongue, and other body parts. It is not known if INGREZZA is safe and effective in children.

Reducing **ING** tardive dyskinesia

ACHIEVING REMISSION

No or minimal movements

at **48 WEEKS** in 59% of people*

WORKS QUICKLY

>5X TD Reduction
at **2 WEEKS** vs placebo†

SIGNIFICANT IMPROVEMENT‡

>10X TD Reduction
at **6 WEEKS** vs placebo§

*In a post-clinical study evaluation of the KINECT 4 trial, 59% (61/103) of people who completed the trial taking INGREZZA (40 mg and 80 mg) achieved a score of 1 or less for each body part measured with AIMS at Week 48. The results are descriptive in nature.

†Reductions based on average change from baseline to Week 2 on an uncontrollable movement severity scale. Results for INGREZZA 80 mg -1.9 (n=77) vs placebo -0.3 (n=76) in a clinical study.

‡INGREZZA was studied in a 6-week clinical trial. A total of 234 people participated in the study. Results were based on 79 people taking the recommended dose of 80 mg.

§Reductions based on average change from baseline to Week 6 on an uncontrollable movement severity scale. Results for INGREZZA 80 mg -3.2 (n=70) vs placebo -0.1 (n=69) in a clinical study.

Select Important Safety Information

Before taking INGREZZA or INGREZZA SPRINKLE, tell your healthcare provider about all of your medical conditions including if you: have liver or heart problems, are pregnant or plan to become pregnant, or are breastfeeding or plan to breastfeed.

Please see additional Important Safety Information on pages 18–19 and accompanying full [Prescribing Information](#), including Boxed Warning, and [Medication Guide](#).



ONCE-DAILY
INGREZZA[®]
(valbenazine) capsules



Actor portrayals

Long-lastING satisfaction

in people taking INGREZZA
for ~2 years*



*55 of 56 patients in a rollover study of patients taking INGREZZA for 48 weeks after completing long-term KINECT 3 & KINECT 4 studies. Based on Patient Satisfaction Questionnaire given to patients at beginning and end of treatment.

Select Important Safety Information

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Make sure you tell all of your healthcare providers that you are taking INGREZZA or INGREZZA SPRINKLE. Taking INGREZZA or INGREZZA SPRINKLE with certain other medicines may cause serious side effects. Especially tell your healthcare provider if you: take digoxin or take or have taken a monoamine oxidase inhibitor (MAOI) medicine. You should not take INGREZZA or INGREZZA SPRINKLE if you are taking, or have stopped taking, a MAOI within the last 14 days.

Please see additional Important Safety Information on pages 18–19 and accompanying full [Prescribing Information](#), including Boxed Warning, and [Medication Guide](#).

ONCE-DAILY
INGREZZA[®]
(valbenazine) capsules

Simple dosING with INGREZZA

INGREZZA makes dosing simple from the start



Starts at an effective dose



Also available in sprinkle formulation



Always one capsule, once daily




Taken any time



With or without food



Can be added to most stable mental health regimens



Not actual size

Actor portrayal


Select Important Safety Information

The most common side effects of INGREZZA or INGREZZA SPRINKLE in people with tardive dyskinesia are sleepiness and tiredness.

Please see additional Important Safety Information on pages 18–19 and accompanying full [Prescribing Information](#), including Boxed Warning, and [Medication Guide](#).



ONCE-DAILY
INGREZZA[®]
(valbenazine) capsules



Not actual size

INGREZZA and INGREZZA® SPRINKLE (valbenazine) capsules is available in 3 effective dosing strengths. The recommended dose of INGREZZA is 80 mg.

Your healthcare provider will start you on 40 mg for 1 week and may keep you on 40 mg, or switch you to 60 mg or 80 mg, depending on your treatment needs.



Have trouble swallowing pills?

There's another way of takING** INGREZZA**

Learn about INGREZZA SPRINKLE at

[INGREZZA.com/TD-Sprinkle](https://www.ingrezza.com/TD-Sprinkle)

Take INGREZZA exactly as your healthcare provider tells you.

Your healthcare provider will tell you what dose of INGREZZA® (valbenazine) capsules to take and when to take it.

- INGREZZA can be taken with or without food
- Do not stop taking INGREZZA without first talking to your healthcare provider
- If you take too much INGREZZA, call your poison control center at 1-800-222-1222

Select Important Safety Information

These are not all of the possible side effects of INGREZZA or INGREZZA SPRINKLE. Call your doctor for medical advice about side effects. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit MedWatch at www.fda.gov/medwatch or call **1-800-FDA-1088**.

Please see additional Important Safety Information on pages 18–19 and accompanying full [Prescribing Information](#), including Boxed Warning, and [Medication Guide](#).



ONCE-DAILY
INGREZZA®
(valbenazine) capsules

With INGREZZA, you can keep **takING** most mental health medicines

In a clinical study, psychiatric status in people taking INGREZZA® (valbenazine) capsules remained stable overall, including those who also took medicine for:



- Depression
- Bipolar disorder
- Schizophrenia/schizoaffective disorder
- Anxiety disorder

Do not stop taking INGREZZA without talking to your healthcare provider first.



“I take INGREZZA once a day, it fits my routine. My TD movements are under control with INGREZZA.”

Individual results may vary

– Moira, living with depression and tardive dyskinesia

Moira was compensated by Neurocrine Biosciences, Inc. to share her story

Select Important Safety Information

Dosage Forms and Strengths: INGREZZA and INGREZZA SPRINKLE are available in 40 mg, 60 mg, and 80 mg capsules.

Please see additional Important Safety Information on pages 18–19 and accompanying full [Prescribing Information](#), including Boxed Warning, and [Medication Guide](#).

ONCE-DAILY
INGREZZA[®]
(valbenazine) capsules



Actor portrayal

People taking INGREZZA for TD can stay on their current dose of most mental health medicines.

Tell your healthcare provider about all the medicines you take, including:



- Prescription medicines
- Over-the-counter medicines
- Vitamins
- Herbal supplements

Select Important Safety Information

INGREZZA or INGREZZA SPRINKLE can cause serious side effects in people with Huntington's disease, including: depression, suicidal thoughts, or suicidal actions. Tell your healthcare provider before you start taking INGREZZA or INGREZZA SPRINKLE if you have Huntington's disease and are depressed (have untreated depression or depression that is not well controlled by medicine) or have suicidal thoughts. Pay close attention to any changes, especially sudden changes, in mood, behaviors, thoughts, or feelings. This is especially important when INGREZZA or INGREZZA SPRINKLE is started and when the dose is changed. Call your healthcare provider right away if you become depressed, have unusual changes in mood or behavior, or have thoughts of hurting yourself.

Please see additional Important Safety Information on pages 18–19 and accompanying full [Prescribing Information](#), including Boxed Warning, and [Medication Guide](#).



ONCE-DAILY
INGREZZA[®]
(valbenazine) capsules

Tak**ING** control

**You've worked hard to improve your mental health.
Don't let uncontrollable movements hold you back.**

Whether your uncontrollable body movements are mild, moderate, or severe, connecting with a psychiatry or neurology healthcare provider who has experience diagnosing and treating TD is important.



Actor portrayals

Select Important Safety Information

Do not take INGREZZA or INGREZZA SPRINKLE if you:

- are allergic to valbenazine, or any of the ingredients in INGREZZA or INGREZZA SPRINKLE.


Please see additional Important Safety Information on pages 18–19 and accompanying full [Prescribing Information](#), including Boxed Warning, and [Medication Guide](#).

ONCE-DAILY
INGREZZA[®]
(valbenazine) capsules



Actor portrayal

Taking control starts with asking your healthcare provider about a treatment for your uncontrollable movements. INGREZZA® (valbenazine) capsules is a treatment proven to reduce TD that's always one capsule, once daily, and #1 prescribed.



Need help starting the conversation about TD? Download our Doctor Discussion Guide at [INGREZZA.com/discussionguide](https://www.ingrezza.com/discussionguide)



Select Important Safety Information

INGREZZA or INGREZZA SPRINKLE can cause serious side effects, including:

- Allergic reactions.** Allergic reactions, including an allergic reaction that causes sudden swelling called angioedema, can happen after taking the first dose or after many doses of INGREZZA or INGREZZA SPRINKLE. Signs and symptoms of allergic reactions and angioedema include: trouble breathing or shortness of breath, swelling of your face, lips, eyelids, tongue, or throat, or other areas of your skin, trouble with swallowing, or rash, including raised, itchy red areas on your skin (hives). Swelling in the throat can be life-threatening and can lead to death. Stop taking INGREZZA or INGREZZA SPRINKLE and go to the nearest emergency room right away if you develop these signs and symptoms of allergic reactions and angioedema.

Please see additional Important Safety Information on pages 18–19 and accompanying full [Prescribing Information](#), including [Boxed Warning](#), and [Medication Guide](#).

ONCE-DAILY
INGREZZA®
 (valbenazine) capsules

Common side effects of INGREZZA

The most common side effects of INGREZZA® (valbenazine) capsules or INGREZZA® SPRINKLE (valbenazine) capsules in people with tardive dyskinesia are sleepiness and tiredness.

INGREZZA or INGREZZA SPRINKLE can cause serious side effects in people with Huntington's Disease, including: depression, suicidal thoughts, or suicidal actions.

What should I do if I experience side effects?

These are not all of the possible side effects of INGREZZA or INGREZZA SPRINKLE. Call your doctor for medical advice about side effects. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit MedWatch at www.fda.gov/medwatch or call **1-800-FDA-1088**.

EmpowerING



Actor portrayals

A lot goes into being the #1 prescribed TD treatment:

- 27 clinical studies with over 1,470 people
- 7 years of real-world experience
- More than 1 million total prescriptions since 2017
- 3 simple and effective dosing options
- Starts at an effective dose



Select Important Safety Information

INGREZZA or INGREZZA SPRINKLE can cause serious side effects, including:

- Sleepiness and tiredness that could cause slow reaction times (somnolence and sedation). Do not drive a car or operate dangerous machinery until you know how INGREZZA or INGREZZA SPRINKLE affects you. Drinking alcohol and taking other medicines may also cause sleepiness during treatment with INGREZZA or INGREZZA SPRINKLE.

Please see additional Important Safety Information on pages 18–19 and accompanying full [Prescribing Information](#), including Boxed Warning, and [Medication Guide](#).

ONCE-DAILY
INGREZZA[®]
(valbenazine) capsules



Actor portrayal

Dedicated to **helpING** you on your treatment journey

The INBRACE® Support Program is designed to help patients who are prescribed INGREZZA® (valbenazine) capsules through a range of financial assistance programs.



Copay Savings Card

If you are eligible, a Savings Card may be applied automatically at the pharmacy, which may help you save money on your INGREZZA prescription.*

*This offer is valid only for patients who have commercial (nongovernment-funded) insurance. Additional terms and conditions apply.

INGREZZA is available through a select network of specialty and local affiliated pharmacies. A specialty pharmacy provides medicines that aren't typically available at your local retail pharmacy.

Once coverage has been authorized, INGREZZA will be shipped directly to you via overnight mail.

Select Important Safety Information

INGREZZA or INGREZZA SPRINKLE can cause serious side effects, including:

- **Heart rhythm problems (QT prolongation).** INGREZZA or INGREZZA SPRINKLE may cause a heart rhythm problem known as QT prolongation. You have a higher chance of getting QT prolongation if you also take certain other medicines during treatment with INGREZZA or INGREZZA SPRINKLE. Tell your healthcare provider right away if you develop any signs or symptoms of QT prolongation, including: fast, slow, or irregular heartbeat (heart palpitations), shortness of breath, dizziness or lightheadedness, or fainting or feeling like you are going to faint.

Please see additional Important Safety Information on pages 18–19 and accompanying full [Prescribing Information](#), including Boxed Warning, and [Medication Guide](#).

ONCE-DAILY
INGREZZA®
(valbenazine) capsules

INGREZZA Start Program

The INGREZZA Start Program is a free trial (one-month supply) of INGREZZA available for new patients.* Your healthcare provider can enroll you in the INGREZZA Start Program by downloading the enrollment form from the INBRACE website and submitting the completed form on your behalf.

INGREZZA Patient Assistance Program

If you do not have prescription coverage for INGREZZA and lack the financial resources to pay for the medicine, you may be able to receive your prescription at no cost through the INGREZZA Patient Assistance Program.† Your healthcare provider can help you apply for the program or call 1-84-INGREZZA to learn more.

*This program is not contingent on a purchase of any kind. Product dispensed under this free trial program may not be submitted for reimbursement to any third-party payer. We reserve the right to modify or cancel the program at any time.

†Additional terms and conditions apply.

To learn more about the INBRACE[®] Support Program,
call 1-84-INGREZZA (1-844-647-3992)
8 AM TO 8 PM ET, Monday through Friday,
or visit [INGREZZA.com/Support](https://www.ingrezza.com/support)

Select Important Safety Information

INGREZZA or INGREZZA SPRINKLE can cause serious side effects, including:

- **Neuroleptic Malignant Syndrome (NMS).** NMS is a serious condition that can lead to death. Call a healthcare provider right away or go to the nearest emergency room if you develop these symptoms and they do not have another obvious cause: high fever, stiff muscles, problems thinking, irregular pulse or blood pressure, increased sweating, or very fast or uneven heartbeat.

Please see additional Important Safety Information on pages 18–19 and accompanying full [Prescribing Information](#), including Boxed Warning, and [Medication Guide](#).



ONCE-DAILY
INGREZZA[®]
(valbenazine) capsules

Important Information

Approved Uses

INGREZZA® (valbenazine) capsules or INGREZZA® SPRINKLE (valbenazine) capsules are prescription medicines used to treat adults with:

- movements in the face, tongue, or other body parts that cannot be controlled (tardive dyskinesia).
- involuntary movements (chorea) of Huntington's disease. INGREZZA or INGREZZA SPRINKLE do not cure the cause of involuntary movements, and do not treat other symptoms of Huntington's disease, such as problems with thinking or emotions.

It is not known if INGREZZA or INGREZZA SPRINKLE is safe and effective in children.

IMPORTANT SAFETY INFORMATION

INGREZZA or INGREZZA SPRINKLE can cause serious side effects in people with Huntington's disease, including: depression, suicidal thoughts, or suicidal actions. Tell your healthcare provider before you start taking INGREZZA or INGREZZA SPRINKLE if you have Huntington's disease and are depressed (have untreated depression or depression that is not well controlled by medicine) or have suicidal thoughts. Pay close attention to any changes, especially sudden changes, in mood, behaviors, thoughts, or feelings. This is especially important when INGREZZA or INGREZZA SPRINKLE is started and when the dose is changed. Call your healthcare provider right away if you become depressed, have unusual changes in mood or behavior, or have thoughts of hurting yourself.

Do not take INGREZZA or INGREZZA SPRINKLE if you:

- are allergic to valbenazine, or any of the ingredients in INGREZZA or INGREZZA SPRINKLE.

INGREZZA or INGREZZA SPRINKLE can cause serious side effects, including:

- **Allergic reactions.** Allergic reactions, including an allergic reaction that causes sudden swelling called angioedema, can happen after taking the first dose or after many doses of INGREZZA or INGREZZA SPRINKLE. Signs and symptoms of allergic reactions and angioedema include: trouble breathing or shortness of breath, swelling of your face, lips, eyelids, tongue, or throat, or other areas of your skin, trouble with swallowing, or rash, including raised, itchy red areas on your skin (hives). Swelling in the throat can be life-threatening and can lead to death. Stop taking INGREZZA or INGREZZA SPRINKLE and go to the nearest emergency room right away if you develop these signs and symptoms of allergic reactions and angioedema.
- **Sleepiness and tiredness that could cause slow reaction times (somnolence and sedation).** Do not drive a car or operate dangerous machinery until you know how INGREZZA or INGREZZA SPRINKLE affects you. Drinking alcohol and taking other medicines may also cause sleepiness during treatment with INGREZZA or INGREZZA SPRINKLE.

- **Heart rhythm problems (QT prolongation).** INGREZZA or INGREZZA SPRINKLE may cause a heart rhythm problem known as QT prolongation. You have a higher chance of getting QT prolongation if you also take certain other medicines during treatment with INGREZZA or INGREZZA SPRINKLE. Tell your healthcare provider right away if you develop any signs or symptoms of QT prolongation, including: fast, slow, or irregular heartbeat (heart palpitations), shortness of breath, dizziness or lightheadedness, or fainting or feeling like you are going to faint.
- **Neuroleptic Malignant Syndrome (NMS).** NMS is a serious condition that can lead to death. Call a healthcare provider right away or go to the nearest emergency room if you develop these symptoms and they do not have another obvious cause: high fever, stiff muscles, problems thinking, irregular pulse or blood pressure, increased sweating, or very fast or uneven heartbeat.
- **Parkinson-like symptoms.** Symptoms include: body stiffness, drooling, trouble moving or walking, trouble keeping your balance, shaking (tremors), or falls.

Before taking INGREZZA or INGREZZA SPRINKLE, tell your healthcare provider about all of your medical conditions including if you: have liver or heart problems, are pregnant or plan to become pregnant, or are breastfeeding or plan to breastfeed.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Make sure you tell all of your healthcare providers that you are taking INGREZZA or INGREZZA SPRINKLE. Taking INGREZZA or INGREZZA SPRINKLE with certain other medicines may cause serious side effects. Especially tell your healthcare provider if you: take digoxin or take or have taken a monoamine oxidase inhibitor (MAOI) medicine. You should not take INGREZZA or INGREZZA SPRINKLE if you are taking, or have stopped taking, a MAOI within the last 14 days.

The most common side effects of INGREZZA or INGREZZA SPRINKLE in people with tardive dyskinesia are sleepiness and tiredness.

The most common side effects of INGREZZA or INGREZZA SPRINKLE in people with chorea associated with Huntington's disease include sleepiness and tiredness, raised itchy red areas on your skin (hives), rash, and trouble getting to sleep or staying asleep.

These are not all of the possible side effects of INGREZZA or INGREZZA SPRINKLE. Call your doctor for medical advice about side effects. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit MedWatch at www.fda.gov/medwatch or call **1-800-FDA-1088**.

Dosage Forms and Strengths: INGREZZA and INGREZZA SPRINKLE are available in 40 mg, 60 mg, and 80 mg capsules.



Treat**ING** your TD movements



INGREZZA® (valbenazine) capsules is proven to reduce TD, always one capsule, once daily, and #1 prescribed.



People taking INGREZZA can stay on most mental health medicines.



Starts at an effective dose.



Take control: start by asking your healthcare provider about INGREZZA.

Scan the QR code or visit [INGREZZA.com/TD](https://www.ingrezza.com/TD) to learn more about treatment.

Select Important Safety Information

INGREZZA or INGREZZA SPRINKLE can cause serious side effects in people with Huntington's disease, including: depression, suicidal thoughts, or suicidal actions. Tell your healthcare provider before you start taking INGREZZA or INGREZZA SPRINKLE if you have Huntington's disease and are depressed (have untreated depression or depression that is not well controlled by medicine) or have suicidal thoughts. Pay close attention to any changes, especially sudden changes, in mood, behaviors, thoughts, or feelings. This is especially important when INGREZZA or INGREZZA SPRINKLE is started and when the dose is changed. Call your healthcare provider right away if you become depressed, have unusual changes in mood or behavior, or have thoughts of hurting yourself.

Please see additional Important Safety Information on pages 18–19 and accompanying full [Prescribing Information](#), including Boxed Warning, and [Medication Guide](#).

